

**UK canine tick-borne
disease in 2016 – an update
References**

References

- Abdullah S, Helps C, Tasker S, Newbury H & Wall R (2016). Ticks infesting domestic dogs in the UK: a large-scale surveillance programme. *Parasites & Vectors* **9**: 39
- Bennett CE (1995) Ticks and Lyme disease. *Advances in Parasitology* **36**: 343–405.
- Bouchard C, Leonard E, Koffi JK et al (2015). The increasing risk of Lyme disease in Canada. *Canadian Veterinary Journal* **56**: 693–9
- Fourie JJ, Stanneck D, Luusa HG, Beugnet F, Wijnveld M et al (2013). Transmission of *Ehrlichia canis* by *Rhipicephalus sanguineus* ticks feeding on dogs and on artificial membranes. *Veterinary Parasitology* **197**:595-603
- Goosens, HAT., Van Den Bogaard, AE. & Nohlmans, MKE (2001). Dogs as sentinels for human Lyme borreliosis in the Netherlands. *Journal of Clinical Microbiology* **39**: 844-8.
- Hansford KM, Pietzsch M, Cull M & Medlock JM (2015). Brown dog tick infestation of a home in England. *Veterinary Record* **176**:129-30
- Holm LP, Kerr MG, Trees AJ, McGarry JM, Munro ER et al (2006). Fatal babesiosis in an untraveled British dog. *Veterinary Record* **159**: 179-80
- Littman MP (2013). Lyme nephritis. *Journal of Veterinary emergency critical care (San Antonio)* **23**:163-73
- Matjila PT, Penzhorn, BL, Bekker CP, Nijhof AM & Jongejan F (2004). Confirmation of occurrence of *Babesia canis vogeli* in domestic dogs in South Africa. *Veterinary Parasitology* **122**: 119–25.
- Phipps L, Del Mar Fernandez De Marco M, Hernández-Triana L, Johnson N, Swainsbury C et al (2016). *Babesia canis* detected in dogs and associated ticks from Essex *Veterinary Record* **178**: 243-4.
- Piesman J (1993). Dynamics of *Borrelia burgdorferi* transmission by nymphal *Ixodes dammini* ticks. *Journal of Infectious Disease* **167**: 1082–5.
- Shaw SE, Binns SH, Birtles RJ, Day MJ, Smithson RC et al (2005). Molecular evidence of tick-transmitted infections in dogs and cats in the United Kingdom. *The Veterinary Record* **157**:645-8
- Shih CM & Spielman A (1993). Accelerated transmission of Lyme disease spirochetes by partially fed vector ticks. *Journal of Clinical Microbiology* **31**: 2878–81.
- Smith F, Ballantyne R, Morgan E & Wall R (2011). The prevalence, distribution and risk associated with tick infestation of dogs in Great Britain. *Medical & Veterinary Entomology* **25**: 377-84.
- Smith FD, Ballantyne R, Morgan E & Wall R (2012). Estimating Lyme disease risk using pet dogs as sentinels. *Comparative Immunology, Microbiology & Infectious Diseases* **35**: 163-7.

Smith FD & Wall LE (2013). Prevalence of Babesia and Anaplasma in ticks infesting dogs in Great Britain. *Veterinary Parasitology* **198**: 18-23

Swainsbury C, Bengtson G & Hill P (2016). Babesiosis in dogs. *Veterinary Record* **178**:172

Wlodarek J, Zuraw A, Walczak R, Dziuban J & Jaskowski JM (2013). Lyme Borreliosis: the most frequent tick-borne disease in humans and animals. *Medycyna Weterynaryjna* **69**: 341-7.

Woodmansey D (2016). New tick-borne disease confirmed in UK dogs *Veterinary Times* **46**: 1